

Lowest Landau level broadened by a Gaussian random potential with an arbitrary correlation length: An efficient continued-fraction approach

Markus Böhm,¹ Kurt Broderix,² and Hajo Leschke¹

¹*Institut für Theoretische Physik, Universität Erlangen-Nürnberg, Staudtstr. 7, D-91058 Erlangen, Germany*

²*Institut für Theoretische Physik, Universität Göttingen, Bunsenstr. 9, D-37073 Göttingen, Germany*

(February 1, 2008)

For an electron in the plane subjected to a perpendicular constant magnetic field and a homogeneous Gaussian random potential with a Gaussian covariance function we approximate the averaged density of states restricted to the lowest Landau level. To this end, we extrapolate the first 9 coefficients of the underlying continued fraction consistently with the coefficients' high-order asymptotics. We thus achieve the first reliable extension of Wegner's exact result [Z. Phys. B **51**, 279 (1983)] for the delta-correlated case to the physically more relevant case of a non-zero correlation length.

Nearly ideal two-dimensional electronic structures have attracted great attention for more than a decade not only because of their varied and important applications, but also because of the discovery of the quantum Hall effect [1]. For a microscopic understanding of the occurring phenomena it is essential to know the spectral properties of electrons confined to two dimensions under the influence of a perpendicular constant magnetic field taking into account the presence of disorder.

A commonly studied minimal model is that of non-interacting electrons which is characterized by the one-electron Hamiltonian given by the Schrödinger operator

$$\hat{H} := \hat{K} + \hat{V} \quad (1)$$

$$\hat{K} := \frac{1}{2m} \left(\frac{\hbar}{i} \frac{\partial}{\partial x_1} - \frac{eB}{2} x_2 \right)^2 + \frac{1}{2m} \left(\frac{\hbar}{i} \frac{\partial}{\partial x_2} + \frac{eB}{2} x_1 \right)^2. \quad (2)$$

Here $x := (x_1, x_2)$ are Cartesian coordinates of the Euclidean plane $\mathbb{R}^2$, $\hbar$ is Planck's constant, e is the elementary charge, m is the (effective) mass of the (spinless) electron, and $B > 0$ the strength of a perpendicular magnetic field. The static random potential V is added to mimic the interaction with quenched disorder. Its probability distribution is assumed to be Gaussian with zero mean and Gaussian covariance, that is

$$\overline{V(x)} = 0, \quad \overline{V(x)V(x')} = \sigma^2 \exp \left[-\frac{(x - x')^2}{2\lambda^2} \right]. \quad (3)$$

The overbar denotes averaging with respect to the probability distribution. Finally, $\sigma > 0$ is the strength and $\lambda > 0$ the correlation length of the fluctuations of the potential.

The spectral resolution of the unperturbed Hamiltonian $\hat{K}$ reads $\hat{K} = \sum_{n=0}^{\infty} \varepsilon_n \hat{E}_n$, where the eigenvalue $\varepsilon_n := (2n+1) \frac{\hbar e B}{2m}$ is called the n -th Landau level and $\hat{E}_n$ denotes the corresponding eigenprojector. With increasing B the magnetic length $l := \sqrt{\hbar/(eB)}$ decreases, while

the degeneracy $\langle x | \hat{E}_n | x \rangle = (2\pi l^2)^{-1}$ (per area) of each Landau level and the distance between successive levels increase. Hence, for a fixed concentration of electrons, sufficiently high fields, and low temperatures it is reasonable to simplify the model by restricting its Hamiltonian (1) to the (still infinite dimensional) eigenspace $\hat{E}_0 L^2(\mathbb{R}^2)$ of the lowest Landau level. This has been done, for example, in the important work of Wegner [5], where the averaged restricted density of states (per area)

$$\varrho(\varepsilon) := \overline{\langle y | \hat{E}_0 \delta(\varepsilon - \hat{E}_0 \hat{H} \hat{E}_0) \hat{E}_0 | y \rangle} \quad (4)$$

has been calculated exactly in the delta-correlated limit $\lambda \downarrow 0$, $\sigma \rightarrow \infty$, $\lambda^2 \sigma^2 = \text{const.}$

The purpose of the present Letter is twofold. One goal is to present what we think is an accurate extension of Wegner's result to arbitrary values of σ and λ . Clearly, this is of physical interest because in the high-field limit the actual correlation length is no longer small in comparison with the magnetic length. The other goal is to demonstrate thereby that the power of the well-known continued-fraction approach to spectral densities of non-trivial quantum problems [2,3] can considerably be enhanced in cases, where one has an *a-priori* information about their asymptotic high-frequency decay.

To get rid of physical dimensions we write ϱ in the standardized form

$$\varrho(\varepsilon) =: \frac{\sqrt{1 + (l/\lambda)^2}}{2\pi l^2 \sigma} W\left(\frac{\lambda^2}{l^2}, \frac{(\varepsilon - \varepsilon_0)}{\sigma} \sqrt{1 + (l/\lambda)^2}\right). \quad (5)$$

In this way we have introduced a one-parameter family of even probability densities on the real line $\mathbb{R}$ with normalized second moment

$$W(a, u) = W(a, -u) \geq 0 \quad (6)$$

$$\int_{\mathbb{R}} du W(a, u) = 1 = \int_{\mathbb{R}} du W(a, u) u^2. \quad (7)$$

Here we have used Eq. (7) of [4] to normalize the second moment.

In this notation Wegner's result for the delta-correlated limit reads [5,6]

$$W(0, u) = \frac{2\pi^{-3/2} \exp(u^2)}{1 + [2\pi^{-1/2} \int_0^u d\xi \exp(\xi^2)]^2}. \quad (8)$$

For the other extreme of the spatial extent of correlations, namely the constantly correlated case $\lambda = \infty$, one simply has [7] the Gaussian

$$W(\infty, u) = (2\pi)^{-1/2} \exp(-u^2/2). \quad (9)$$

For intermediate values of a no exact expression for $W(a, u)$ is known. What is exactly known, however, is

the fact that $W(a, u)$ falls off for sufficiently large $|u|$ like a Gaussian. More precisely, by Eqs. (17) and (15) of [4] (see also [8])

$$\lim_{u \rightarrow \pm\infty} \frac{1}{u^2} \ln(W(a, u)) = -\frac{a+2}{2a+2}. \quad (10)$$

In the sequel we will design approximations to the Stieltjes transform

$$R(a, z) := \int_{\mathbb{R}} du \frac{W(a, u)}{z - iu}, \quad \operatorname{Re} z > 0, \quad (11)$$

of the standardized density of states W which in turn will yield approximations to W by means of the inversion formula

$$W(a, u) = \frac{1}{\pi} \lim_{v \downarrow 0} \operatorname{Re} [R(a, v + iu)]. \quad (12)$$

According to Stieltjes' classical theory, see for example [9,10], R can be expanded into a Jacobi-type continued fraction

$$R(a, z) = \underset{j=1}{\overset{\infty}{\text{K}}} \left(\frac{r_j(a)}{z} \right), \quad r_j(a) \geq 0. \quad (13)$$

Here we are using the notation

$$\underset{j=1}{\overset{\infty}{\text{K}}} \left(\frac{\Delta_j}{z} \right) := \frac{1}{z + \frac{\Delta_1}{z + \frac{\Delta_2}{z + \ddots}}} \quad (14)$$

for the continued fraction with coefficients $\Delta_1, \Delta_2, \dots$ and variable z .

To derive the continued-fraction coefficients $\{r_j(0)\}$ and $\{r_j(\infty)\}$ corresponding to (8) and (9), respectively, we use the identity

$$\underset{j=1}{\overset{\infty}{\text{K}}} \left(\frac{\beta + \gamma j}{z} \right) = \frac{D_{-(\beta/\gamma)-1}(\gamma^{-1/2}z)}{\gamma^{1/2} D_{-\beta/\gamma}(\gamma^{-1/2}z)} =: T(\beta, \gamma, z), \quad (15)$$

valid if $\gamma > 0$, $\beta + \gamma > 0$, and $\operatorname{Re} z > 0$. Here D_ν denotes Whittaker's parabolic cylinder function with index ν , see Section 8.1 of [11]. The identity follows by iteration from the observation that the rhs of (15) obeys

$$T(\beta, \gamma, z) = [z + (\beta + \gamma)T(\beta + \gamma, \gamma, z)]^{-1} \quad (16)$$

which itself is a consequence of the first of the four recurrence relations for the D_ν 's in Section 8.1.3 of [11]. In fact, Eq. (15) is equivalent to Eq. (14) in §50 of [9]. With the help of (12) one now checks that

$$r_j(0) = \frac{1}{2} + \frac{1}{2}j, \quad r_j(\infty) = j, \quad (17)$$

for all $j \geq 1$.

According to [12] the asymptotic behavior (10) implies the following asymptotic linear growth for the continued-fraction coefficients

$$\lim_{j \rightarrow \infty} \frac{r_j(a)}{j} = \frac{a+1}{a+2}. \quad (18)$$

It is natural to match this linear growth to the first $J < \infty$ coefficients $r_1(a), \dots, r_J(a)$ to construct the announced approximations $R^{(J)}(a, z)$ to $R(a, z)$ by means of

$$R^{(J)}(a, z) := \mathbb{K}_{j=1}^{\infty} \left(\frac{r_j^{(J)}(a)}{z} \right), \quad (19)$$

where

$$r_j^{(J)}(a) := \begin{cases} r_j(a) & \text{for } j \leq J \\ r_J(a) + \frac{a+1}{a+2}(j-J) & \text{for } j > J \end{cases}. \quad (20)$$

Not surprisingly, the approximation $R^{(J)}$ to R results in an approximation $W^{(J)}$ to W with the same Gaussian fall-off (10) in the tails. This can be seen as follows. By virtue of (15) the approximation $R^{(J)}$ can be expressed as a terminating continued fraction

$$R^{(J)}(a, z) = \frac{1}{z + \frac{r_1(a)}{z + \frac{r_2(a)}{z + \dots + \frac{r_{J-1}(a)}{z + \frac{r_J(a)}{z + r_J(a)T(r_J(a), \frac{a+1}{a+2}, z)}}}}}. \quad (21)$$

In view of (12) it is therefore sufficient to show

$$\lim_{u \rightarrow \pm\infty} \frac{1}{u^2} \ln(\operatorname{Re}[T(\beta, \gamma, iu)]) = -\frac{1}{2\gamma} \quad (22)$$

and

$$\lim_{u \rightarrow \pm\infty} \operatorname{Im}[T(\beta, \gamma, iu)] = 0, \quad (23)$$

because these asymptotic properties are conserved under the (repeated) substitution

$$T(\beta, \gamma, iu) \mapsto [iu + \alpha T(\beta, \gamma, iu)]^{-1} \quad (24)$$

for any $\alpha > 0$. The validity of (22) und (23) can be deduced from an asymptotic evaluation [13] of the following Riccati differential equation

$$\gamma \frac{d}{du} T(\beta, \gamma, iu) = i\beta [T(\beta, \gamma, iu)]^2 - uT(\beta, \gamma, iu) - i \quad (25)$$

which itself follows from the aforementioned recurrence relations. To summarize, Eq. (15) adds a two-parameter family of terminators for affine-linear extrapolations to the toolkit of [2].

In order to compute the first J continued-fraction coefficients $r_1(a), \dots, r_J(a)$, we employ the well-known one-to-one correspondence [9,10,2] to the first J even moments $M_2(a), \dots, M_{2J}(a)$ of the standardized density of states.

According to (4) and (5) one has for the latter

$$\begin{aligned} M_{2j}(\lambda^2/l^2) &:= \int_{\mathbb{R}} du W(\lambda^2/l^2, u) u^{2j} \\ &= 2\pi l^2 \left(\frac{1 + l^2/\lambda^2}{\sigma^2} \right)^j \overline{\langle y | (\hat{E}_0 \hat{V} \hat{E}_0)^{2j} | y \rangle} \end{aligned} \quad (26)$$

if $j \geq 1$. We now substitute $\hat{V} = \int_{\mathbb{R}^2} d^2x V(x) |x\rangle \langle x|$ into the rhs of (26) and use the standard reduction formula

$$\overline{\prod_{k=1}^{2j} V(x(k))} = \sum_{k=1}^{(2j-1)!!} \prod_{s=1}^j \overline{V(x(P_k(2s-1))) V(x(P_k(2s)))} \quad (27)$$

for the average of a product of $2j$ jointly Gaussian random variables with zero mean. Here $x(1), \dots, x(2j) \in \mathbb{R}^2$ are $2j$ points of the plane and P_k denotes the k -th of the $(2j-1)!!$ permutations of the first $2j$ natural numbers which lead to different sets $\{\{P_k(1), P_k(2)\}, \dots, \{P_k(2j-1), P_k(2j)\}\}$ of j pairs $\{P_k(2s-1), P_k(2s)\}$. Since the covariance function (3) of the random potential V and the position representation

$$\langle x | \hat{E}_0 | x' \rangle = \frac{1}{2\pi l^2} \exp \left[\frac{i}{2l^2} (x_1 x'_2 - x_2 x'_1) - \frac{1}{4l^2} (x - x')^2 \right] \quad (28)$$

of the eigenprojector corresponding to the lowest Landau level are both Gaussian, one ends up with a sum over 4j-dimensional Gaussian integrals which can be performed to yield

$$M_{2j}(a) = \left(1 + \frac{1}{a}\right)^j \sum_{k=1}^{(2j-1)!!} \frac{1}{\det(A_k(a))}, \quad a > 0. \quad (29)$$

Here the $2j \times 2j$ -matrix $A_k(a)$ is defined through its entries in terms of the Kronecker delta

$$(A_k(a))_{\mu, \nu} := \left(1 + \frac{1}{a}\right) \delta_{\mu, \nu} - \delta_{\mu+1, \nu} - \frac{1}{a} \sum_{s=1}^j (\delta_{\mu, P_k(2s-1)} \delta_{\nu, P_k(2s)} + \delta_{\nu, P_k(2s-1)} \delta_{\mu, P_k(2s)}). \quad (30)$$

We have computed the sum (29) for $j = 1, 2, \dots, 6$ using the AXIOM symbolic system [14] for general a . The results for $M_{2j}(a)$, $j = 1, 2, 3, 4$, are given by the following rational functions

$$M_2(b-1) = 1$$

$$M_4(b-1) = (3b^2 + 2)/(b^2 + 1)$$

$$M_6(b-1) = (15b^6 + 75b^4 + 102b^2 + 30)/(b^6 + 6b^4 + 11b^2 + 6)$$

$$M_8(b-1) = (105b^{24} + 2835b^{22} + 32830b^{20} + 213697b^{18} + 861130b^{16} + 2231807b^{14} + 3750338b^{12} + 4038543b^{10} + 2717117b^8 + 1105510b^6 + 261216b^4 + 32792b^2 + 1680)/$$

$$(b^{24} + 29b^{22} + 365b^{20} + 2620b^{18} + 11854b^{16} + 35276b^{14} + 69974b^{12} + 91906b^{10} + 78025b^8 + 41015b^6 + 12461b^4 + 1954b^2 + 120).$$

(31)

Since the expressions for $M_{10}(a)$ and $M_{12}(a)$ are extremely lengthy, we only give, as an example, their values for $a = 1$:

$$M_{10}(1) = \frac{158\,659\,605\,940\,126\,452\,841}{294\,310\,802\,651\,335\,470}$$

$$M_{12}(1) = \frac{388\,336\,271\,072\,847\,928\,549\,926\,597\,113\,071\,401\,088\,677\,997\,478\,405\,727\,555\,223\,031}{82\,363\,680\,790\,265\,452\,914\,044\,225\,729\,941\,466\,953\,642\,191\,484\,142\,275\,602\,750}.$$

(32)

The moments $M_{14}(a)$ and $M_{16}(a)$ have been computed exactly as a reduced fraction for $a = \frac{1}{4}, \frac{1}{2}, 1, 2, 4$ only, by employing a C-program. For the same set of values for a we also have computed $M_{18}(a)$, but only have been able to gather up the $17!! \approx 3.4 \cdot 10^7$ terms occurring in (29) to a sum of approximately 10^4 reduced fractions, the value of which has been computed accurately by floating-point arithmetics. For lack of space we only show the first 35 digits for the example $a = 1$:

$$M_{14}(1) = 47657.946072630475536554559639613945 \dots$$

$$M_{16}(1) = 545841.43592501744224295351679205436 \dots$$

$$M_{18}(1) = 6980770.3795705620571551127542699202 \dots$$

From Fig. 1 it can be seen that the resulting continued-fraction coefficients $r_1(a), \dots, r_9(a)$ approach quite rapidly their asymptotic behavior given in (18). For example, the relative extrapolation error $|r_9^{(8)}(a) - r_9(a)|/r_9(a)$ varies between 0.2% for $a = 4$ and 0.02% for $a = \frac{1}{4}$.

In Fig. 2 we demonstrate the convergence of the resulting approximations $W^{(J)}(1, u)$ to $W(1, u)$ for increasing $J = 1, 2, \dots, 9$. The differences of two successive approximations, $W^{(J)}(a, u) - W^{(J-1)}(a, u)$, seem to form a nearly alternating sequence with geometric decrease for fixed u . Both observations, taken together, suggest rapid point-wise convergence of the sequence $\{W^{(J)}\}$. Therefore, we may safely conclude that $W^{(9)}$ constitutes a reliable approximation to W . Figure 3 shows a plot of the corresponding approximation $\varrho^{(9)}$ to the averaged density of states ϱ for different λ . Note that, by construction, $\varrho^{(9)}$ is exact in the limiting cases of a delta-correlated and a constantly correlated random potential; more importantly, $\varepsilon \mapsto 2\pi l^2 \varrho^{(9)}(\varepsilon + \varepsilon_0)$ is an even probability density with the same first 18 moments and the same Gaussian fall-off in the tails as the true density for general values of the correlation length λ .

-
- [1] For review-type literature see T. Ando, A. B. Fowler, and F. Stern, *Rev. Mod. Phys.* **54**, 437 (1982); I. V. Kukushkin, S. V. Meshkov, and V. B. Timofeev, *Usp. Fiz. Nauk.* **155**, 219 (1988) [*Sov. Phys. Usp.* **31**, 511 (1988)]; G. Landwehr (ed), *High Magnetic Fields in Semiconductor Physics I, II, III* (Springer, Berlin, 1987, 1989, 1992). M. Janßen, O. Viehweger, U. Fastenrath, and J. Hajdu, *Introduction to the Theory of the Integer Quantum Hall Effect* (VCH, Weinheim, 1994)
 - [2] V. S. Viswanath and G. Müller, *The Recursion Method — Application to Many-Body Dynamics*, Lecture Notes in Physics, Vol m23 (Springer, Berlin, 1994), and references therein.
 - [3] Continued fractions were applied successfully to a lattice version of (1) by G. Czycholl and W. Ponischowski, *Z. Phys. B* **73**, 343 (1988).
 - [4] K. Broderix, N. Heldt, and H. Leschke, *J. Phys. A* **24**, L825 (1991).
 - [5] F. Wegner, *Z. Phys. B* **51**, 279 (1983).
 - [6] see also E. Brézin, D. J. Gross, and C. Itzykson, *Nucl. Phys. B* **235**, 24 (1984); A. Klein and J. F. Perez, *Nucl. Phys. B* **251**, 199 (1985).
 - [7] see, for example, K. Broderix, H. Heldt, and H. Leschke, *Z. Phys. B* **68**, 19 (1987).
 - [8] W. Apel, *J. Phys. C* **20**, L577 (1987).
 - [9] O. Perron, *Die Lehre von den Kettenbrüchen, Vols. I, II* (Teubner, Stuttgart, 1977).
 - [10] H. S. Wall, *Continued Fractions* (Van Nostrand, New York, 1948)
 - [11] W. Magnus, F. Oberhettinger, and R. P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, 3rd enlarged ed. (Springer, Berlin, 1966).
 - [12] D. S. Lubinsky, H. N. Mhaskar, and E. B. Saff, *Constr. Approx.* **4**, 65 (1988), and references therein.
 - [13] see, for example, C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers*, (McGraw-Hill, Auckland, 1978)
 - [14] R. D. Jenks and R. S. Sutor, *AXIOM: The Scientific Computation System* (Springer, Berlin, 1992).




FIG. 1. Symbols represent the first 9 continued-fraction coefficients $r_1(a), \dots, r_9(a)$ for different values of a . Note that $r_1(a) = 1$ for all a by (7). The straight lines correspond to the asymptotics of $\{r_j^{(9)}(a)\}$ as defined in (20).




FIG. 2. Difference of two successive approximations to W for $a=1$ as a function of u given on a linear (a) and a logarithmic (b) scale.




FIG. 3. The approximation $\rho^{(9)}$ to the averaged density of states ρ for different values of the correlation length λ as a function of the energy ε .